

Lampiran 13. Hasil uji anova nilai R/C ratio atas biaya total menurut klasifikasi irigasi dan musim tanam tahun 2012

Notes

| | | |
|------------------------|--------------------------------|--|
| Output Created | 13-Nov-2013 08:24:10 | |
| Comments | | |
| Input | Data | C:\Users\user\Documents\eny\data uji beda RC ratio.sav |
| | Active Dataset | DataSet1 |
| | Filter | <none> |
| | Weight | <none> |
| | Split File | <none> |
| | N of Rows in Working Data File | 160 |
| Missing Value Handling | Definition of Missing | User-defined missing values are treated as missing. |
| | Cases Used | Statistics for each analysis are based on cases with no missing data for any variable in the analysis. |
| Syntax | | ONEWAY RCratio BY klasifikasi /STATISTICS DESCRIPTIVES HOMOGENEITY /MISSING ANALYSIS /POSTHOC=SCHEFFE ALPHA(0.05). |
| Resources | Processor Time | 00:00:00.032 |
| | Elapsed Time | 00:00:00.022 |

| | N | Mean | Std. Deviation | Std. Error | 95% Confidence Interval for Mean | | Minimum | Maximum |
|-------|-----|--------|----------------|------------|----------------------------------|-------------|---------|---------|
| | | | | | Lower Bound | Upper Bound | | |
| 1 | 40 | 2.7705 | .43605 | .06895 | 2.6310 | 2.9100 | 1.28 | 3.38 |
| 2 | 40 | 1.9068 | .30293 | .04790 | 1.8099 | 2.0036 | .67 | 2.40 |
| 3 | 40 | 2.1010 | .43085 | .06812 | 1.9632 | 2.2388 | 1.39 | 3.28 |
| 4 | 40 | 1.3975 | .28208 | .04460 | 1.3073 | 1.4877 | .93 | 2.01 |
| Total | 160 | 2.0439 | .61456 | .04859 | 1.9480 | 2.1399 | .67 | 3.38 |

Test of Homogeneity of Variances

RCratio

| Levene Statistic | df1 | df2 | Sig. |
|------------------|-----|-----|------|
| 2.814 | 3 | 156 | .041 |

ANOVA

RCratio

| | Sum of Squares | df | Mean Square | F | Sig. |
|----------------|----------------|-----|-------------|--------|------|
| Between Groups | 38.714 | 3 | 12.905 | 94.348 | .000 |
| Within Groups | 21.337 | 156 | .137 | | |
| Total | 60.051 | 159 | | | |

Post Hoc Tests

Multiple Comparisons

RCratio

Scheffe

| (I) klasifikasi | (J) klasifikasi | Mean Difference (I-J) | Std. Error | Sig. | 95% Confidence Interval | |
|-----------------|-----------------|-----------------------|------------|------|-------------------------|-------------|
| | | | | | Lower Bound | Upper Bound |
| 1 | 2 | .86375* | .08270 | .000 | .6300 | 1.0975 |
| | 3 | .66950* | .08270 | .000 | .4358 | .9032 |
| | 4 | 1.37300* | .08270 | .000 | 1.1393 | 1.6067 |
| 2 | 1 | -.86375* | .08270 | .000 | -1.0975 | -.6300 |
| | 3 | -.19425 | .08270 | .142 | -.4280 | .0395 |
| | 4 | .50925* | .08270 | .000 | .2755 | .7430 |
| 3 | 1 | -.66950* | .08270 | .000 | -.9032 | -.4358 |
| | 2 | .19425 | .08270 | .142 | -.0395 | .4280 |
| | 4 | .70350* | .08270 | .000 | .4698 | .9372 |
| 4 | 1 | -1.37300* | .08270 | .000 | -1.6067 | -1.1393 |
| | 2 | -.50925* | .08270 | .000 | -.7430 | -.2755 |
| | 3 | -.70350* | .08270 | .000 | -.9372 | -.4698 |

*. The mean difference is significant at the 0.05 level.

Homogeneous Subsets

RCratio

Scheffe

| klasifikasi | N | Subset for alpha = 0.05 | | |
|-------------|----|-------------------------|--------|--------|
| | | 1 | 2 | 3 |
| 4 | 40 | 1.3975 | | |
| 2 | 40 | | 1.9068 | |
| 3 | 40 | | 2.1010 | |
| 1 | 40 | | | 2.7705 |
| Sig. | | 1.000 | .142 | 1.000 |

Means for groups in homogeneous subsets are displayed.