

Appendix 12.

Table of Frequency

pretest

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	30.00	1	3.1	3.1	3.1
	40.00	5	15.6	15.6	18.8
	45.00	6	18.8	18.8	37.5
	50.00	8	25.0	25.0	62.5
	55.00	5	15.6	15.6	78.1
	60.00	5	15.6	15.6	93.8
	65.00	2	6.3	6.3	100.0
	Total	32	100.0	100.0	

posttest

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	70.00	3	9.4	9.4	9.4
	75.00	4	12.5	12.5	21.9
	80.00	6	18.8	18.8	40.6
	85.00	7	21.9	21.9	62.5
	90.00	7	21.9	21.9	84.4
	95.00	4	12.5	12.5	96.9
	100.00	1	3.1	3.1	100.0
	Total	32	100.0	100.0	

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent

pretest	32	100.0%	0	.0%	32	100.0%
posttest	32	100.0%	0	.0%	32	100.0%

Descriptives

			Statistic	Std. Error
pretest	Mean		50.1563	1.46334
	95% Confidence Interval for Mean	Lower Bound	47.1718	
		Upper Bound	53.1407	
	5% Trimmed Mean		50.2431	
	Median		50.0000	
	Variance		68.523	
	Std. Deviation		8.27787	
	Minimum		30.00	
	Maximum		65.00	
	Range		35.00	
	Interquartile Range		10.00	
	Skewness		-.144	.414
	Kurtosis		-.234	.809
posttest	Mean		84.2188	1.42184
	95% Confidence Interval for Mean	Lower Bound	81.3189	
		Upper Bound	87.1186	
	5% Trimmed Mean		84.2361	
	Median		85.0000	
	Variance		64.693	
	Std. Deviation		8.04317	
	Minimum		70.00	
	Maximum		100.00	

Range	30.00	
Interquartile Range	10.00	
Skewness	-.126	.414
Kurtosis	-.719	.809

Notes

Output Created	16-Mar-2013 13:21:38	
Comments		
Input	Data	D:\Neng_eRa\Titipan\T-Test Bagus Dona.sav
	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	32
Missing Value Handling	Definition of Missing	User-defined missing values for dependent variables are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any dependent variable or factor used.
Syntax	EXAMINE VARIABLES=pretest posttest /PLOT BOXPLOT STEMLEAF /COMPARE GROUPS /STATISTICS DESCRIPTIVES /INTERVAL 95 /MISSING LISTWISE /NOTOTAL.	
Resources	Processor Time	00:00:01.092
	Elapsed Time	00:00:01.495

Histogram

